


## Convex Hull



6/8/2002 12:42 PM

Convex Hull

1

## Outline and Reading

- ◆ Convex hull (§12.5.2)
- ◆ Orientation (§12.5.1-2)
- ◆ Sorting by angle (§12.5.5)
- ◆ Graham scan (§12.5.5)
- ◆ Analysis (§12.5.5)


6/8/2002 12:42 PM

Convex Hull


2

## Convex Polygon

- ◆ A convex polygon is a nonintersecting polygon whose internal angles are all convex (i.e., less than  $\pi$ )
- ◆ In a convex polygon, a segment joining two vertices of the polygon lies entirely inside the polygon



convex



nonconvex


6/8/2002 12:42 PM

Convex Hull

3

## Convex Hull

- ◆ The convex hull of a set of points is the smallest convex polygon containing the points
- ◆ Think of a rubber band snapping around the points



6/8/2002 12:42 PM

Convex Hull

4

## Special Cases

- ◆ The convex hull is a segment
  - Two points
  - All the points are collinear
- ◆ The convex hull is a point
  - there is one point
  - All the points are coincident




6/8/2002 12:42 PM

Convex Hull

5

## Applications

- ◆ Motion planning
  - Find an optimal route that avoids obstacles for a robot
- ◆ Geometric algorithms
  - Convex hull is like a two-dimensional sorting




6/8/2002 12:42 PM

Convex Hull

6

# Computing the Convex Hull

- The following method computes the convex hull of a set of points
  - Phase 1:** Find the lowest point (anchor point)
  - Phase 2:** Form a nonintersecting polygon by sorting the points counterclockwise around the anchor point
  - Phase 3:** While the polygon has a nonconvex vertex, remove it



6/8/2002 12:42 PM

Convex Hull

7


# Orientation

- The orientation of three points in the plane is clockwise, counterclockwise, or collinear

- orientation**( $a, b, c$ )
  - clockwise (CW, right turn)
  - counterclockwise (CCW, left turn)
  - collinear (COLL, no turn)

- The orientation of three points is characterized by the sign of the determinant  $\Delta(a, b, c)$ , whose absolute value is twice the area of the triangle with vertices  $a, b$  and  $c$

$$\Delta(a, b, c) = \begin{vmatrix} x_a & y_a & 1 \\ x_b & y_b & 1 \\ x_c & y_c & 1 \end{vmatrix}$$




6/8/2002 12:42 PM

Convex Hull

8

# Sorting by Angle

- Computing angles from coordinates is complex and leads to numerical inaccuracy
- We can sort a set of points by angle with respect to the anchor point  $a$  using a comparator based on the orientation function
  - $b < c \Leftrightarrow \text{orientation}(a, b, c) = \text{CCW}$
  - $b = c \Leftrightarrow \text{orientation}(a, b, c) = \text{COLL}$
  - $b > c \Leftrightarrow \text{orientation}(a, b, c) = \text{CW}$




6/8/2002 12:42 PM

Convex Hull

9

# Removing Nonconvex Vertices

- Testing whether a vertex is convex can be done using the orientation function
- Let  $p, q$  and  $r$  be three consecutive vertices of a polygon, in counterclockwise order
  - $q$  convex  $\Leftrightarrow \text{orientation}(p, q, r) = \text{CCW}$
  - $q$  nonconvex  $\Leftrightarrow \text{orientation}(p, q, r) = \text{CW or COLL}$



6/8/2002 12:42 PM


Convex Hull

10

# Graham Scan

- The Graham scan is a systematic procedure for removing nonconvex vertices from a polygon
- The polygon is traversed counterclockwise and a sequence  $H$  of vertices is maintained

for each vertex  $r$  of the polygon  
 Let  $q$  and  $p$  be the last and second last vertex of  $H$   
**while**  $\text{orientation}(p, q, r) = \text{CW or COLL}$   
 remove  $q$  from  $H$   
 $q \leftarrow p$   
 $p \leftarrow$  vertex preceding  $p$  in  $H$   
 Add  $r$  to the end of  $H$



6/8/2002 12:42 PM

Convex Hull

11

# Analysis

- Computing the convex hull of a set of points takes  $O(n \log n)$  time
  - Finding the anchor point takes  $O(n)$  time
  - Sorting the points counterclockwise around the anchor point takes  $O(n \log n)$  time
    - Use the orientation comparator and any sorting algorithm that runs in  $O(n \log n)$  time (e.g., heap-sort or merge-sort)
  - The Graham scan takes  $O(n)$  time
    - Each point is inserted once in sequence  $H$
    - Each vertex is removed at most once from sequence  $H$

6/8/2002 12:42 PM

Convex Hull

12